REVIEW ARTICLE

UPI JOURNAL OF CHEMICAL AND LIFE SCIENCES (UPI-JCLS)

ISSN: 2581-4648 (An International online Peer Reviewed, Open Access Journal)

www.uniquepubinternational.com Published by Unique Pub International (UPI)

SCHIFF BASE CONDENSATION VERSUS ALTERNATIVE PREPARATIVE ROUTES: A COMPARATIVE REVIEW OF REACTION EFFICIENCY, PRODUCT QUALITY AND ENVIRONMENTAL COST

RAGHAD MOHAMMED SALIH

Department of Chemistry, College of Education, University of Al-Qadisiyah, Diwaniyah, Iraq

***Corresponding Author**
Raghad Mohammed Salih
E-Mail ID: raghad.mohammed@qu.edu.iq

Received: 03 08 2026 Revised: 21 08 2026 Accepted: 26 09 2026

Abstract
Schiff base condensation, the reaction of a primary amine with an aldehyde or a ketone to give an azomethine, is still one of the most heavily used preparative operations in coordination, medicinal and materials chemistry. Its appeal rests on a deceptively simple picture: two components, one new bond, water as the only by-product. In practice the way that bond is made varies enormously, and the choice between the procedures available is seldom argued on quantitative grounds. This review compares the classical reflux preparation of Schiff bases with the non-conventional routes that have largely displaced it, namely microwave irradiation, ultrasonic irradiation, mechanochemical grinding and milling, catalysis by natural acids, ionic liquids and deep eutectic solvents, and aqueous or one-pot protocols. The imine route is then placed beside other strategies that deliver comparable targets: Mannich aminomethylation, diazotisation-coupling, reductive amination, and the copper-catalysedazide-alkyne cycloaddition. Reported times, isolated yields, catalyst demand and the few available green metrics are collected from primary reports and compared directly, along with the direct-versus-template question governing how the metal complexes are assembled. Three points recur. Energy-input methods compress reaction times by one to three orders of magnitude while usually raising yield by ten to twenty percentage points; on atom economy the condensation is matched only by the Mannich reaction and by catalytic reductive amination, and is far ahead of diazotisation-coupling and amide coupling; and the reversibility that makes the imine easy to form remains its practical weakness. A minimum reporting set is proposed to make future comparisons meaningful.

Keywords: *Schiff base; azomethine; condensation reaction; green synthesis; microwave and ultrasonic irradiation mechanochemistry; template synthesis; comparative preparative methods.*

Copyright:© 2026 The author(s). This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

INTRODUCTION

Few reactions in the synthetic repertoire have been reused as often, and with as little change, as the condensation Hugo Schiff described in the middle of the nineteenth century. The azomethine group it produces, -CH=N-, is small, easy to install and unusually informative: it carries a lone pair suitable for coordination, a polarised double bond that responds to substituents, and a diagnostic infrared band that makes the reaction easy to follow. That combination explains why Schiff bases and their metal complexes continue to appear across such different fields, from antimicrobial and anticancer screening [1] to corrosion protection [2], ion sensing [3], asymmetric catalysis [4] and the construction of crystalline porous frameworks [5].

What has changed is not the reaction but the way it is run. For most of the twentieth century the standard operation was refluxing equimolar carbonyl compound and amine in ethanol or methanol, usually with a drop of acetic acid, for anything between an hour and a full day [6]. Over the last two decades that procedure has been progressively displaced, in the published record at least, by microwave irradiation, sonication, solventless grinding and catalysis with materials as ordinary as lemon juice [7]. Reviews of these developments are numerous and several are good [8, 9, 10], but almost all treat the greener methods as self-evidently superior and stop there.

The more useful question, and the one addressed here, runs in two directions at once. How do the various ways of making a Schiff base compare when time, isolated yield, purification effort and waste are set side by side? And how does the condensation itself compare with the other coupling strategies a chemist might reasonably choose for a related

target: Mannich aminomethylation, diazotisation-coupling, reductive amination, or click cycloaddition? Mass-based green metrics answer part of both questions [11], but they are still rarely reported here. This review assembles what has been reported, compares it as carefully as the primary data allow, and marks where the comparison cannot yet be made.

2. WHAT ACTUALLY GOVERNS THE CONDENSATION

2.1 A two-step mechanism with a narrow optimum

Although textbooks draw imine formation as a single arrow, it is two distinct events separated by a change in rate-determining step (Figure 1). The amine adds to the carbonyl carbon to give a tetrahedral carbinolamine, and that intermediate then loses water to give the C=N bond. Classical kinetic work on aliphatic systems established both stages and, importantly,




Figure 01: The Schiff base condensation drawn as the two elementary steps that it actually comprises: nucleophilic addition of the amine to the carbonyl carbon to give a carbinolamine, followed by acid-catalysed dehydration to the azomethine. The reaction is reversible, and the water it releases drives the reverse process. showed that acid catalyses the second but suppresses the first by protonating the amine [12, 13]. The consequence is the familiar bell-shaped pH-rate profile shown in Figure 2, with a maximum that for a weakly basic aromatic amine sits somewhere between pH 4 and 5 and moves upward as the amine becomes more basic [14]. Above that region dehydration becomes rate-limiting; below it, the concentration of free amine collapses and the addition step controls the rate.




Figure 02: Dependence of the rate of imine formation on pH for three amine basicities. The curves are constructed from the classical two-step rate law and are illustrative only; they are not experimental measurements. Below the maximum the concentration of free amine limits the rate, above it dehydration of the carbinolamine does, and the maximum therefore moves to higher pH as the amine becomes more basic.

Subsequent investigations refined rather than overturned this image. Polarographic measurement of the benzaldehyde-hydrazine system made it possible to separate the formation and dehydration equilibria of the carbinolamine and estimate its protonation constants [15] and equilibrium work on formaldehyde with weakly basic anilines showed how small the formation constants become when the nucleophile is poor [16]. Steric and hydrogen bonding factors could

change the profile further [17], sometimes resulting in many breaks in the curve. Later, computations in explicit water reproduced the essential features and made the role of the solvent explicit: water molecules relieve the ring strain of the proton-transfer transition states, and acid catalysis provides much of the driving force for the forward reaction [18, 19]. The practical lesson to the bench chemist is trite but often overlooked. A catalytic amount of a weak acid is helpful; a stoichiometric amount is not.

2.2 Substrate electronics and the water problem

Aromatic aldehydes with electron withdrawing groups condense faster and form more stable products than those with electron donating groups. Ketones are always more challenging than aldehydes for the obvious steric and electronic reasons. Weakly basic aromatic amines are also slow partners and with them the equilibrium constant itself might become so small that the imine is impossible to isolate at all [20]. The kinetic and thermodynamic selectivity might be opposed when there is competition of many amines for one carbonyl molecule. This is exactly the characteristic that is utilised in dynamic covalent libraries [21].

The reaction produces water, and the water can drive the reverse reaction. Practically every preparative variation to be examined below can be viewed as a different answer to the same question: how do you get rid of water, or make its presence irrelevant? Azeotropic removal, molecular sieves, solventless circumstances, quick heating to alter the equilibrium before hydrolysis can compete, template capture of the product by a metal ion, all remedies to that one problem. Table 01 shows the structural classes generated by the same condensation depending on the choice of the carbonyl and amine partners.

Table 01: Principal structural classes of Schiff bases, with the carbonyl and amine partners from which each is obtained.

Class	Carbonyl partner	Amine partner	Characteristic feature
Aldimine (benzylidene amine)	Aromatic or aliphatic aldehyde	Primary amine	The common case; -CH=N-, usually crystalline and easy to purify
Ketimine	Ketone	Primary amine	Slower to form, more readily hydrolysed, often needs forcing conditions
Salicylaldimine	2-Hydroxybenzaldehyde	Primary amine	Intramolecular O-H...N hydrogen bond; strong N,O chelation
Salen type (N2O2)	Salicylaldehyde (2 equiv.)	Diamine (1 equiv.)	Tetradentate; the reference ligand class in asymmetric catalysis
Acyclic bis-Schiff base	Dialdehyde, or 2 equiv. aldehyde	2 equiv. amine, or 1 equiv. diamine	Two azomethine units joined by a flexible spacer
Macrocyclic ([1+1], [2+2])	Dialdehyde	Diamine	Ring closure; usually requires a metal template to give one product
Hydrazone	Aldehyde or ketone	Hydrazine or acyl hydrazide	C=N-N unit; markedly more resistant to hydrolysis
Thiosemicarbazone	Aldehyde or ketone	Thiosemicarbazide	N,N,S donor set; heavily studied for biological activity
Oxime	Aldehyde or ketone	Hydroxylamine	C=N-OH; much larger formation constant than a simple imine
Amino-acid Schiff base	Salicylaldehyde	alpha-Amino acid	Water soluble; biologically relevant model system
Polymer-supported Schiff base	Aldehyde	Free -NH2 of chitosan or a polyamine	Grafting of the azomethine onto a macromolecular support
Imine-linked framework	Polytopic aldehyde	Polytopic amine	Crystalline porous network; reversibility is essential to crystallisation

3. THE CONVENTIONAL PREPARATION, AND WHERE IT BECOMES EXPENSIVE

The normal procedure is easy to say. One equiv. of carbonyl compound and 1 equiv. of amine are mixed in an alcohol, a few drops of glacial acetic acid added and the mixture refluxed until analysis by thin-layer chromatography shows the limiting reagent to be exhausted. Times recorded for this surgery vary greatly. Ortho-hydroxyketimines derived from hindered ketones were refluxed at 140 degrees Celsius for at least twenty-four hours [22]. Imidazole derived systems were refluxed for about an hour before the reaction was judged complete [23]. In a systematic study of eight dithiocarbazate and thiosemicarbazone Schiff bases the conventional route was the slowest of the four procedures examined [24]. "This has three costs. The first is energy, the hours of reflux at an alcohol's boiling point repeated across a sequence of twelve or twenty analogues is a huge and largely unmeasured expense. The second is a solvent. Recrystallisation requires 20-30 ml ethanol for a typical 1 mmol preparation, a similar amount is required for the reaction, and solvent dominates the waste inventory, but does not appear in the reaction equation. The third expense is less visible and it is about the product, not the method. The heating for a prolonged period in a protic solvent leads to partial hydrolysis, and in the case of sensitive substrates side reactions, due to the water produced by the reaction. In several comparative studies it is reported that the conventionally prepared samples required further purification while the non-conventional prepared samples did not [24, 25]. This does not render the classical method wrong. It remains the most reliable route if a substrate is thermally resilient, if the scale is big and if no specialised equipment is available, and is still the method against which every alternative is measured [26].

4. ENERGY-INPUT ROUTES TO THE SAME IMINE

4.1 Microwave irradiation

Of the alternatives, microwave-assisted synthesis has been studied most and is the most convincing. Dielectric heating raises the bulk temperature of a polar mixture far faster than an oil bath can, and because the imine forms quickly once the barrier is crossed, the time saving is dramatic rather than marginal. The comparison reported by Kundu and co-workers is representative: reactions that took twenty-four hours under reflux were complete in one minute under microwave irradiation, and the isolated yields rose from 79-87% to 94-97% [22]. Imidazole Schiff bases show the same behaviour from the other side of the comparison: 90-98% yield within two to four minutes under irradiation, against roughly an hour at reflux [23].

The method is not restricted to simple benzylidene anilines. Calix[4]resorcinarene Schiff bases were obtained in 80-91% yield under mild microwave conditions in ethanol [27], pyrimidine-based congeners in 72-81% [28], and catalyst-free protocols on aminophenols give phenolic Schiff bases directly, with a work-up that amounts to little more than washing [29]. Natural catalysts combine well with the technique: cashew shell extract used as a natural acid under solventless microwave conditions gave one-pot access to a series of Schiff bases with high yield and, according to the authors, elimination of side products [30]. Attempts have also been made to model the process quantitatively, relating yield to applied power and molar ratio [31]. Metal complexes have been prepared the same way, either from the isolated ligand or in a single operation [32, 33], and recent surveys of the field are broadly consistent in their conclusions [34, 35].

4.2 Ultrasonic irradiation

Sonication works by a different physical mechanism. Acoustic cavitation produces transient bubbles whose collapse generates local hot spots and intense micro-mixing, so the benefit appears mainly in heterogeneous or poorly mixed systems rather than through bulk heating. For Schiff base chemistry the practical outcome is similar to microwave activation, though usually less extreme: reaction times of minutes instead of hours, and yields improved by a modest but consistent margin. In one of the few three-way comparisons available, a catalyst-free series of bis-Schiff bases prepared from aromatic aldehydes and diamines was run under conventional heating, microwave irradiation and ultrasound, and the sonochemical route gave the products in high yield and the shortest times of the three [36]. Ultrasonic preparation of Schiff bases from 4-aminoantipyrine in ethanol with glacial acetic acid reached 86% within five minutes [37], and 2-aminopyridine derivatives condensed with 3-ethoxysalicylaldehyde rapidly enough that the products could be grown as single crystals and characterisedcrystallographically [38].

Where sonication is genuinely distinctive is in the preparation of the metal complexes. Because nucleation is fast and mixing is efficient, sonochemically prepared complexes are frequently obtained as nanostructured solids with narrow size distributions, which is difficult to achieve by slow reflux [39, 40]. Nano-sized Fe(II), Cd(II) and Zn(II) complexes prepared this way have been used directly as precursors to metal oxide nanoparticles [41], and Ag(I) and Ni(II) complexes obtained sonochemically performed well as visible-light photocatalysts [42]. An elegant recent example assembles manganese bis(imino)pyridine complexes in one pot and one step in thirty minutes, without an added acid catalyst and without any attempt to exclude water, replacing what had been a multiday, multistep sequence [43]. Direct comparisons of the ultrasonic and conventional routes for the same ligand and the same complexes confirm the pattern [44, 45].

4.3 Grinding and ball milling

Mechanochemistry removes the solvent altogether. In its simplest form the two solids are ground together in a mortar for a few minutes; in its instrumented form a vibrational or planetary mill does the same work reproducibly and at scale [46, 47]. Fluorinated imines prepared by manual grinding of equimolar substrates at room temperature were obtained in good to excellent yields after only fifteen minutes, and the authors note that these yields were comparable to or

higher than those reported for the same compounds by other methods [48]. Chromone-derived Schiff bases were obtained in 80-81% isolated yield by the same approach [49], and imine formation proceeds well even in the solid state on a polymer backbone, where efficient energy delivery by ball collisions promotes rapid condensation without any chemical solvent [50].

The strength of the approach is its waste profile rather than its speed, and this is where the few quantitative assessments are most useful: systematic evaluation of mechanochemical protocols against solution chemistry, using atom economy, E-factor, process mass intensity and related indicators, shows a consistently lower environmental footprint, driven mainly by the absence of bulk solvent and by better stoichiometric control [51, 52]. Its limitation is equally clear. Liquid or low-melting substrates are awkward, scale-up of manual grinding is not credible, and a small amount of liquid additive is often needed anyway, which weakens the solvent-free claim. A three-way comparison of microwave, ultrasonic and mechanochemical preparation of quinazolinone Schiff bases found yields spanning 11-90% across the series, a reminder that the mechanochemical option is substrate-sensitive in a way the others are not [53].

4.4 Catalysts and reaction media

A parallel line of work leaves the heating conventional and changes the catalyst instead. Fruit juices containing citric, ascorbic and malic acids have been used as natural acid catalysts under solvent-free conditions, giving Schiff bases in good yields with a work-up limited to filtration and washing [54]. Yields of this kind are fully competitive with mineral acid catalysis: condensation of vanillin with p-aminoacetophenone under lime juice catalysis, by grinding alone, gave 94.45% of the isolated imine [55]. The approach extends to multicomponent condensations run in water [56] and to substrates as demanding as benzoxazolinones [57]. The literature on lemon juice as a general biocatalyst and medium is now substantial [58], and the same chemistry has been adapted for undergraduate teaching, where the pedagogical value of a reaction run in citrus juice is obvious [59].

Designer solvents occupy the other end of the same spectrum. Ionic liquids can act as solvent and catalyst simultaneously, and an imidazolium system used with microwave heating gave pyrrolopyrimidine Schiff bases in 82-94% yield with good recovery of the medium [60]. Deep eutectic solvents offer much of the same behaviour at a fraction of the cost and with better biodegradability [61, 62, 63]. Water-tolerant Lewis acids such as metal triflates deserve particular mention: at low loading they accelerate imine formation and exchange enough to give near-quantitative yields of crystalline imine-linked networks under mild conditions [64]. Heterogeneous oxide catalysts extend the reaction upstream, generating the aldehyde in situ from an alcohol so that the imine is formed in a single operation from a more accessible feedstock [65]; a related strategy reduces a nitroarene in situ and condenses the resulting amine under milling [66]. Broader surveys of metal-free and bio-based methodology place these developments in context [67].

4.5 Water, one-pot assemblies and continuous operation

Running the condensation in water is counter-intuitive, since water is the by-product, but it works more often than expected. Salicylaldehyde-derived Schiff bases were obtained in excellent yield in water under microwave irradiation, whereas the same substrates under thermal conditions gave lower yields and a more tedious work-up [25]. Imine-linked frameworks have likewise been assembled in water in high yield, and under microwave conditions the reaction time falls considerably [68]. Reversibility can even be turned into a synthetic tool: using imine exchange rather than de novo condensation allowed a crystalline framework to be prepared by simple stirring under reflux in twenty minutes [69]. One-pot multicomponent variants, in which the Schiff base is generated and consumed without isolation, are common in heterocyclic synthesis [70] and in the direct assembly of complexes [71].

Table 02: Representative preparations of Schiff bases and their metal complexes, grouped by activation route. Values are those stated in the primary reports; n.r. indicates that the source did not report the quantity.

System	Route	Conditions and reaction time	Isolated yield (%)	Ref.
ortho-Hydroxyketimines (12 compounds)	Conventional	Reflux at 140 °C, at least 24 h	79-87	[22]
ortho-Hydroxyketimines (12 compounds)	Microwave, solvent-free	1 min	94-97	[22]
Imidazole Schiff bases	Conventional	Reflux in ethanol, about 1 h	n.r.	[23]
Imidazole Schiff bases	Microwave	Ethanol, 2-4 min	90-98	[23]
Salicylaldehyde + aromatic amines	Conventional	Thermal, in water; tedious work-up	lower than MW (n.r.)	[25]
Salicylaldehyde + aromatic amines	Microwave	Water as solvent, one step	excellent (n.r.)	[25]
Pyrimidine Schiff base congeners	Microwave	Two microwave steps, 300 W	72-81	[28]

System	Route	Conditions and reaction time	Isolated yield (%)	Ref.
Calix[4]resorcinarene Schiff bases	Microwave	Ethanol, mild conditions	80-91	[27]
Benzylidene (phenolic) Schiff bases	Microwave, catalyst-free	Solventless	high (n.r.)	[29]
Schiff bases with cashew shell extract	Microwave, natural acid	Solvent-free, one pot	high (n.r.)	[30]
Pyrrolo[2,3-d]pyrimidine Schiff bases	Microwave + ionic liquid	[HMIM][TFSI] as solvent and catalyst	82-94	[60]
Schiff base of o-vanillin and its complexes	Microwave vs conventional	Both routes run for comparison	n.r.	[32]
Cr(III) Schiff base and mixed-ligand complexes	Microwave vs conventional	Shorter time and lower energy by MW	n.r.	[33]
Bis-Schiff bases from diamines	Conventional / MW / ultrasound	All three compared, no catalyst	high by US (n.r.)	[36]
4-Aminoantipyrine Schiff bases (3a-k)	Ultrasound	40 kHz, ethanol, AcOH, 0-5 min	up to 86	[37]
2-Aminopyridine Schiff bases	Ultrasound	Short irradiation; crystals obtained	high (n.r.)	[38]
Mn bis(imino)pyridine complexes	Ultrasound	One pot, one step, 30 min, methanol	n.r.	[43]
Ferrocenyl Schiff base and complexes	Ultrasonication-solvothermal	One-pot equimolar condensation	high mass yield (n.r.)	[45]
Fluorinated imines (12 new compounds)	Manual grinding	Solvent-free, room temperature, 15 min	good to excellent (n.r.)	[48]
Chromone-derived imines	Grinding	Solvent-free	80-81	[49]
Vanillin + p-aminoacetophenone	Grinding, lime juice catalyst	Solvent-free	94.45	[55]
Quinazolinone Schiff bases (27 compounds)	MW / US / milling	Ethanol; no further purification needed	11-90	[53]
Amino-acid Schiff bases and Cu(II) complexes	Conventional vs pestle and mortar	Reflux in ethanol vs solid state	n.r.	[26]
Dithiocarbazate and thiosemicarbazone Schiff bases (8)	Conventional vs grinding, water, lemon juice	All four compared	n.r.	[24]
Schiff base and Ni(II) complex from dichloroaniline	Conventional vs green	Green route faster and higher yielding	n.r.	[82]

5. LIGAND FIRST OR COMPLEX FIRST: DIRECT VERSUS TEMPLATE PREPARATION

For metal complexes the choice of preparation is not only about energy input. It is about order of operations, and the two possibilities are set out in Figure 3. In the stepwise route the Schiff base is isolated, purified and characterised, and only then treated with the metal salt. In the template route the metal ion is present while the condensation occurs, and it organises the fragments around itself. The template effect was recognised early in macrocyclic chemistry and has been exploited ever since [72, 73]. Its value is greatest when the target is a ring. Condensation of a dialdehyde with a diamine in the absence of a metal ion generally gives a mixture of [1+1], [2+2] and [3+3] products that is difficult to separate; in the presence of a suitable cation, one of these is selected and isolated cleanly [74]. The selection is genuinely sensitive to the metal: ionic radius, coordination number and preferred geometry, and even the halide counter-ion, determine which macrocycle forms [75]. In some systems the metal stops the condensation halfway, giving a half-condensed ligand

that cannot be obtained otherwise [76], and in others the choice of cation decides between a tridentate and a tetradentate salen-type product [77]. Reviews of pyridine-containing macrocycles and of one-pot macrocyclic complexes document how general this control has become [78, 79].




Figure 03: Stepwise and template routes to a Schiff base metal complex. In the stepwise route the ligand is isolated and characterised before metallation; in the template route the metal ion selects one macrocycle from what would otherwise be a mixture of ring sizes.

The template route has a well-known cost: the metal is difficult to remove afterwards, so the free ligand may be inaccessible [80]. Non-template preparations avoid that problem and give a ligand that can be fully characterised before complexation, which matters when the ligand itself is the object of study [81]. Combining a template with an energy-input method is increasingly common, and ultrasonication combined with solvothermal treatment has been used to reach ferrocenyl Schiff base complexes in high yield [45]. In short, the two routes answer different questions, and reporting a template preparation as though it were simply a faster way to the same compound is misleading.

6. PREVIOUS STUDIES: READING THE COMPARATIVE LITERATURE TOGETHER

A number of groups have done the useful thing and run the same substrates by two or more methods in the same laboratory, sometimes extending the comparison to the derived metal complexes [82]. Their results are collected in Table 2 and summarised by activation route in Figure 4; read together they support several statements that no single paper establishes on its own. The first is that the time saving is real and large. Across the comparisons in Table 2 the reduction in reaction time is rarely less than one order of magnitude and is sometimes three, the extreme case being twenty-four hours reduced to one minute [22]. The second is modest gains in yields. Typically the improvement is a gain of ten to twenty percentage points, and is biggest where the conventional yield was poor. This is consistent with the view that quick heating outruns breakdown and hydrolysis rather than affecting the inherent thermodynamics. Third, the purity is also usually increased; some authors indicate that the product obtained by the non-traditional technique was pure enough to be used without recrystallisation, whereas the conventional product was not [25, 53]. Fourth, in the case of three green methods being compared with each other instead of with reflux, the most common ranking places microwave first, ultrasound second and grinding or natural acid catalysis third [7, 83], albeit with a narrow margin and clearly substrate-dependent. Two warnings need to be made. Very few of these studies disclose energy use and practically none indicate solvent mass therefore the environmental claim is mainly based on the qualitative reasoning that less time and less solvent must mean less impact. Also, the comparisons are not always fair: conventional conditions are occasionally taken from an older article instead of re-optimised, which flatters the new method. Where re-optimisation has been done properly, the advantage of the non-conventional route is smaller but still present [8]. Table 3 sets the four strategies against each other feature by feature.




Figure 04: Isolated yields reported for Schiff base preparations, grouped by the route used to drive the condensation. Ranges are those given in the primary reports collected in Table 2; a single marker denotes a single reported value.

Table 03: The four ways of driving the condensation, compared feature by feature. Entries summarise the studies collected in Table 02.

Feature	Conventional reflux	Microwave irradiation	Ultrasonic irradiation	Grinding or ball milling
Typical reaction time	1-24 h	seconds to a few minutes	2-30 min	5-30 min
Typical isolated yield	70-90%	80-98%	80-95%	80-95%
Solvent demand	20-30 mL per mmol, plus recrystallisation	little or none	a few mL, mainly for cavitation	none, or a drop of grinding liquid
Equipment	standard glassware	dedicated or domestic microwave reactor	ultrasonic bath or probe	mortar and pestle, or a mixer or planetary mill
Scale usually demonstrated	up to gram and above	1-5 mmol	1-5 mmol	1-5 mmol, gram scale reported
Solid product obtained	well formed crystals	often improved crystallinity	small, narrowly distributed particles	microcrystalline or partly amorphous
Principal limitation	time, energy and solvent	poor coupling for non-polar mixtures	limited benefit in homogeneous solution	strongly substrate-dependent; liquids awkward

7. THE SCHIFF BASE ROUTE SET AGAINST OTHER PREPARATION METHODS

So far the comparison has been internal. It is more interesting to ask how the condensation compares with the other reactions a chemist might use to attach a nitrogen-containing fragment to an aromatic scaffold, since the choice of reaction usually decides the shape of the whole project.

7.1 Mannich aminomethylation

The Mannich reaction condenses an active-hydrogen compound, formaldehyde and a secondary amine to give a beta-amino carbonyl compound, and the resulting Mannich bases occupy much the same biological territory as Schiff bases [84]. The two reactions are often used in sequence, with a triazole Schiff base aminomethylated to give the corresponding Mannich base [85]. As a preparative operation the Mannich reaction is more demanding: three

components rather than two, formaldehyde as an unpleasant reagent, regiochemical questions when the substrate has more than one acidic position, and a product linked through a flexible methylene bridge rather than a rigid, conjugated C=N unit [86]. Its compensating advantages are that the product is not hydrolytically labile in the way an imine is, and that the amine component can be varied almost without limit [87].

7.2 Diazotisation and coupling

Azo compounds are made by diazotising a primary aromatic amine at 0-5 degrees Celsius and coupling the diazonium salt with an electron-rich partner [88]. The chromophore obtained is far stronger than an azomethine, which is why azo chemistry dominates dye and colorimetric-reagent work. The preparative cost, however, is high. The reaction needs strict temperature control, strong mineral acid, and a nitrite salt; the diazonium intermediate is unstable and in some cases hazardous; and the yields reported for well-behaved systems, 45-80% in a representative series, are lower than those routinely obtained for imines [89]. Efforts to green the sequence, for example by replacing mineral acid with an acidic ion-exchange resin, address some of these problems but not the low-temperature requirement [90]. Recent surveys of azo synthesis describe the same migration towards microwave, ultrasonic and solvent-free protocols seen in Schiff base chemistry [91].

7.3 Reductive amination and amide coupling

If the target is a stable secondary amine rather than an imine, reductive amination is the obvious route, and mechanistically it simply adds a reduction step to the same condensation [92]. Catalytic versions using molecular hydrogen are now highly developed [93, 94], and enzymatic reductive aminases carry out condensation and reduction in one active site with excellent enantioselectivity [95]. The trade-offs are straightforward. The product is robust and configurationally defined, but the route needs a reductant, a catalyst, and often pressure equipment, and selectivity between primary, secondary and tertiary products is a genuine difficulty. Amide coupling, the other common way to join an amine to a carbon fragment, is more reliable still but is notorious for its poor atom economy because of the coupling reagents involved.

7.4 Azide-alkyne cycloaddition

The copper-catalysedazide-alkyne cycloaddition is the modern benchmark for reliability. It is high-yielding, regioselective, tolerant of water and of most functional groups, and gives a triazole that is essentially inert to hydrolysis, oxidation and reduction [96, 97]. Against a Schiff base condensation it wins decisively on product stability and on functional-group tolerance. It loses on three counts: it needs a metal catalyst, whose removal is a real problem in biological applications and has driven much of the work on heterogeneous copper systems [98]; it needs an organic azide, which restricts substrate availability and raises safety concerns that only recent reagent development has begun to address [99]; and it requires both partners to be pre-functionalised, so the apparent one-step simplicity conceals two preparative steps upstream [100].

7.5 The cost side: green metrics

Placed on the mass-based metrics, and using the model reactions collected in Table 4, the imine condensation sits at the top of the table (Figure 5). It joins two molecules and expels one water molecule, so its theoretical atom economy for a typical benzylidene aniline is close to 91%, the only loss being the eighteen mass units of water. The Mannich reaction is almost identical in principle, though it carries a third component. Catalytic reductive amination with hydrogen is marginally higher still, at about 91%, since the added mass is only two units; with borohydride it falls to roughly 88%. Amide coupling with a carbodiimide is far worse, because the urea by-product is heavier than the fragment it delivered. The azide-alkyne cycloaddition is the interesting case: the cycloaddition itself is fully atom-economical, but preparing the azide from a benzylic halide brings the two-step figure down to about 70%, and the catalyst, ligand and solvent are counted nowhere in any of these numbers [11, 101].

Atom economy alone, however, is a poor guide, as several authors have argued forcefully [102]. The E-factor, which counts everything, usually tells a different story, because for small-scale preparations solvent dominates the waste inventory regardless of the reaction chosen. This is exactly why the solventless and aqueous variants of the Schiff base condensation matter more than their yield improvements suggest. Where the full set of mass and energy metrics has been applied to a comparable condensation, the reductions achieved by switching to a green protocol have been large: in one detailed treatment of azachalcone synthesis, waste production fell by roughly 95% on the E-factor and the energy intensity dropped by about two orders of magnitude [103]. Comparable analyses of multicomponent routes to pharmaceuticals reach similar conclusions [104].




Figure 05: Atom economy calculated for the model reactions listed in Table 4. The values follow from the stoichiometry alone and take no account of solvent, catalyst, work-up or the preparation of the coupling partners themselves.

Table 04: Model reactions used for the atom-economy comparison in Figure 5. Values are calculated from the stoichiometry shown and are not taken from any experimental report.

Route	Model reaction	By-products counted	Atom economy (%)
Schiff base condensation	PhCHO + PhNH2 ->PhCH=NPh	H2O	90.96
Mannich aminomethylation	PhCOCH3 + HCHO + HN(CH3)2 -> PhCOCH2CH2N(CH3)2	H2O	90.77
Reductive amination (H2)	PhCHO + PhNH2 + H2 -> PhCH2NHPh	H2O	91.05
Reductive amination (NaBH4)	PhCHO + PhNH2 + 0.25 NaBH4 -> PhCH2NHPh	borate salts	87.80
Azide-alkyne cycloaddition (cycloaddition step only)	PhCH2N3 + PhC(triple bond)CH -> 1-benzyl-4-phenyl-1,2,3-triazole	none	100.00
Azide-alkyne route counted from the halide	PhCH2Br + NaN3, then PhC(triple bond)CH	NaBr	69.57
Diazotisation and coupling	PhNH2 + NaNO2 + 2 HCl + PhOH -> 4-hydroxyazobenzene	2 H2O, NaCl, HCl	60.22
Amide coupling with DCC	PhCOOH + PhNH2 + DCC ->PhCONHPh	dicyclohexylurea	46.78

8. PRODUCT QUALITY: PURITY, CRYSTALLINITY AND PARTICLE SIZE

The preparation route affects more than the number in the yield column. Because non-conventional methods reach completion before side reactions accumulate, the crude products are frequently cleaner, and several authors report that recrystallisation could be omitted entirely [53]. The solid-state properties differ too. Sonochemical preparation, with its rapid and spatially uniform nucleation, tends to give small, narrowly distributed particles, and this has been used deliberately to obtain nanoscale complexes [39, 40]. Microwave preparation has been reported to improve crystallinity in complexes prepared under otherwise identical conditions [34]. Mechanochemical products, by contrast, are often microcrystalline or partly amorphous, which is an advantage for dissolution but a disadvantage if single crystals are required for structure determination. These differences are rarely discussed, yet they matter directly for applications where surface area or particle size governs performance [105].

9. VERIFYING THAT THE IMINE IS REALLY THERE

The analytical burden is light because the sole objective of the reaction is one new bond, but it has to be satisfied properly. The primary evidence is the disappearance of the carbonyl stretch around 1700 cm-1 and the appearance of an azomethine band in the region of between about 1580 and 1650 cm-1, sometimes higher for strongly conjugated systems. This is the observation reported in essentially every paper cited here [49]. This is confirmed by proton NMR which shows a singlet for the azomethine at about 8.3-8.6 ppm and the absence of the aldehyde proton [55]. Mass spectrometry and elemental analysis determine the composition and a proportion of the literature single-crystal diffraction settles the geometry and supports the E configuration [38].

Table 5 collects the observations that together make the case. For metal complexes the diagnostic set expands. A shift of the azomethine band to lower wavenumber on complexation indicates coordination through the imine nitrogen, molar conductance distinguishes electrolytes from non-electrolytes, magnetic moments and electronic spectra suggest the geometry, and thermogravimetry identifies lattice and coordinated water [41, 105]. What is often missing, and should not be, is any evidence that the product prepared by the new method is the same substance as that prepared by the old one. Melting point and infrared agreement is the minimum; a mixed melting point or an overlaid diffraction pattern would be better.

Table 05: Analytical evidence normally used to establish that the azomethine bond has formed and, for the complexes, that it coordinates.

Observation	What it establishes	Behaviour usually reported
Loss of the carbonyl stretch	The aldehyde or ketone has been consumed	Band near 1700 cm-1 disappears
New azomethine stretch	The C=N bond has formed	Roughly 1580-1650 cm-1; higher for strongly conjugated systems
Azomethine singlet in 1H NMR	One imine proton per condensation	Near 8.3-8.6 ppm; the CHO signal near 9.8-10.0 ppm is absent
Imine carbon in 13C NMR	Confirms the C=N carbon	Commonly 155-165 ppm
Molecular ion in MS	Molecular formula	M+ or [M+H]+ consistent with the loss of one H2O
Elemental analysis	Composition	Found within about 0.4% of calculated
Single-crystal diffraction	Geometry and configuration	E configuration about C=N; planarity of the chelate ring
Shift of the azomethine band on complexation	Coordination through the imine nitrogen	Shift of roughly 10-30 cm-1 to lower wavenumber
Molar conductance	Electrolytic or non-electrolytic complex	Distinguishes ionic from neutral species
Magnetic moment and electronic spectra	Oxidation state and geometry	Square planar, tetrahedral or octahedral assignment
Thermogravimetry	Lattice and coordinated water	Stepwise mass loss below and above about 150 °C
Powder diffraction	Crystallinity and phase purity	Sharp reflections for crystalline solids; halo for amorphous ones
Comparison with the conventionally prepared sample	That the two routes give the same substance	Agreement of melting point, infrared and NMR; rarely reported

10. REVERSIBILITY: THE SAME PROPERTY THAT HELPS AND HURTS

The imine bond is formed under thermodynamic control, and it can be broken the same way. In synthesis this is an advantage, because errors are corrected during assembly; it is the basis of dynamic covalent chemistry and the reason imine-linked frameworks crystallise at all [5, 106]. It also makes possible tricks that are impossible with irreversible bonds, such as preparing a framework by exchange in twenty minutes [69] or depolymerising and reconstructing it at the end of its life. In application the same property is a liability. Imines hydrolyse in aqueous acid, and the concern has been raised explicitly in the corrosion-inhibition literature, where a compound is asked to survive in 1 M hydrochloric acid for hours; one group addressed it directly by monitoring the imine absorption band to demonstrate that their

ligand remained intact [107]. Metal ions can also promote hydrolysis rather than protect against it, and complexes have been isolated containing both the intact ligand and its hydrolysis product in the same molecule [108]. The strategies developed for frameworks are instructive: strong interlayer hydrogen bonding that shields the C=N from water [109], or chemical locking of the imine into an irreversible heterocycle [110, 111]. Reviews of imine-linked frameworks treat this as the central design problem of the field [112].

11. HOW THE CHOSEN ROUTE FOLLOWS THE COMPOUND INTO ITS APPLICATION

The preparation is not indifferent to what the compound is later required to do. Regarding corrosion inhibition, efficiencies over 90% in 1 M hydrochloric acid have now become commonplace for Schiff bases with several heteroatoms and extended conjugated systems [113, 114, 115] and the same molecules can be prepared by thermal or microwave condensation with no reported difference in performance [116]. Inhibitors based on chitosan, obtained by condensation of an aldehyde with the free amino groups of the biopolymer, exemplify the direct transfer of the process to macromolecular substrates [117].

In sensing, where limits of detection are often in the nm range, the ligand must be spectroscopically pure, therefore purification is more important than speed [118, 119]. The salen family is the reference case in catalysis and its commercial use in hydrolytic kinetic resolution gives a view of what a well-behaved Schiff base ligand may support [120, 121]. In each of these sectors the path is selected for a different reason. This is precisely the argument against treating any one strategy as universally optimum.

12. GAPS, WEAKNESSES AND WHAT SHOULD BE REPORTED

Three weaknesses in the current literature stand out. Reporting is inconsistent: reaction time is almost always given, isolated yield usually, solvent volume rarely, and energy consumption almost never, so the environmental claims that justify the newer methods cannot generally be checked. Comparisons are often unfair, with re-optimised new conditions set against literature values for the old ones. And scale is nearly always small; the great majority of the microwave, ultrasonic and grinding preparations discussed here were carried out on 1-5 mmol, which tells us little about whether the advantage survives a hundredfold increase, although flow and milling equipment now exists that would allow the question to be answered [51].

Three opportunities follow. The first is straightforward: report the numbers. Table 6 sets out a minimum reporting set that would cost an author nothing and would make cross-laboratory comparison possible. The second is to treat the substrate scope honestly, since the substrate-sensitivity of mechanochemistry in particular is under-reported. The third is to close the loop between preparation and property, by preparing the same compound by two routes and testing both, rather than assuming that identical spectra imply identical performance. Figure 6 summarises, as a practical map, how the choice might be made in the meantime.




Figure 06: A practical map for choosing a preparative route, based on what the target is and on how the substrate behaves rather than on which method is currently fashionable.

Table 06: Minimum reporting set proposed for comparative preparative studies of Schiff bases, with an indication of how often each item is currently given.

Item to report	Why it matters	How often reported at present
Scale, in mmol of limiting reagent	Yield and time are scale dependent	Usually
Total solvent volume, reaction and work-up	Solvent dominates the waste inventory	Rarely
Catalyst identity, loading and recovery	Determines whether the route is catalytic in practice	Sometimes
Nominal power and measured energy input	The only basis for an energy claim	Almost never
Time to full conversion, by TLC or HPLC	Distinguishes conversion from isolated yield	Usually, but the method is often unstated
Isolated yield after a stated purification	Makes yields comparable between laboratories	Usually
Purity, and the method used to measure it	Separates a clean product from a fast one	Sometimes
Direct comparison with the conventional sample	Confirms the two routes give the same compound	Rarely
E-factor or process mass intensity	Turns a green claim into a number	Almost never
Number of repeats and the spread of yields	Shows whether the advantage is reproducible	Almost never

CONCLUSION

Schiff base condensation remains the most economical way to install a nitrogen-carbon double bond, and nothing in the comparative literature challenges that position. What has changed is how the reaction is run. Microwave irradiation, sonication, mechanochemical grinding and catalysis by inexpensive natural acids all compress reaction times from hours to minutes or seconds, generally raise isolated yields by ten to twenty percentage points, and reduce or remove the solvent that dominates the waste inventory. Ranked against the alternatives, the condensation is as atom-economical as catalytic reductive amination and the Mannich reaction, far ahead of diazotisation-coupling and amide coupling, and ahead of the azide-alkyne route once the preparation of the azide is counted; it is simpler to run than any of them, and less stable than all of them. For metal complexes the decision is a different one, between a template assembly that can select a single macrocycle and a stepwise route that yields a fully characterised free ligand. The unresolved issue is not chemical but evidential. Until reaction mass, solvent volume and energy are reported alongside yield and time, the environmental case for the greener protocols will rest on plausibility rather than measurement.

FUNDING

Nil

INFORM CONSENT AND ETHICAL STATEMENT

Not Applicable

ACKNOWLEDGEMENT

Not Declared

REFERENCES

1. Abd-El-Aziz, A.; Li, Z.; Zhang, X.; Elnagdy, S.; Mansour, M. S.; ElSherif, A.; Ma, N.; Abd-El-Aziz, A. S. Advances in Coordination Chemistry of Schiff Base Complexes: A Journey from Nanoarchitectonic Design to Biomedical Applications. Topics in Current Chemistry 2025, 383, 20. DOI: 10.1007/s41061-025-00489-w.
2. Hosseini, M.; Mertens, S. F. L.; Ghorbani, M.; Arshadi, M. R. Asymmetrical Schiff bases as inhibitors of mild steel corrosion in sulphuric acid media. Materials Chemistry and Physics 2003, 78, 800-808. DOI: 10.1016/S0254-0584(02)00390-5.

3. Khan, S.; Chen, X.; Almahri, A.; Allehyani, E. S.; Alhumaydhi, F. A.; Ibrahim, M. M.; Ali, S. Recent developments in fluorescent and colorimetric chemosensors based on Schiff bases for metallic cations detection: A review. Journal of Environmental Chemical Engineering 2021, 9, 106381. DOI: 10.1016/j.jece.2021.106381.
4. Shaw, S.; White, J. D. Asymmetric Catalysis Using Chiral Salen-Metal Complexes: Recent Advances. Chemical Reviews 2019, 119, 9381-9426. DOI: 10.1021/acs.chemrev.9b00074.
5. Segura, J. L.; Mancheno, M. J.; Zamora, F. Covalent organic frameworks based on Schiff-base chemistry: synthesis, properties and potential applications. Chemical Society Reviews 2016, 45, 5635-5671. DOI: 10.1039/C5CS00878F.
6. Odularu, A. T. Ease to Challenges in Achieving Successful Synthesized Schiff Base, Chirality, and Application as Antibacterial Agent. BioMed Research International 2023, 2023, 1626488. DOI: 10.1155/2023/1626488.
7. Mahmood, A. A. J. Green synthesis of Schiff bases: a review study. Iraqi Journal of Pharmacy 2022, 18, 180-193. DOI: 10.33899/iphr.2022.170406.
8. Ismaeel, M. I.; Parveen, B.; Dogar, S. S.; Aftab, K.; Abbas, K.; Munawar, K. S. Comparing green and conventional methods for Schiff base synthesis and unveiling environmental stability applications: a review. Journal of Coordination Chemistry 2024, 77, 921-959. DOI: 10.1080/00958972.2024.2362344.
9. Kolodziej, B. Comparison of conventional and green approaches to the synthesis of aromatic Schiff bases. Polish Journal of Chemical Technology 2024, 26, 39-55. DOI: 10.2478/pjct-2024-0039.
10. Thakur, S.; Bhalla, A. Sustainable synthetic endeavors of pharmaceutically active Schiff bases and their metal complexes: A review on recent reports. Tetrahedron 2024, 153, 133836. DOI: 10.1016/j.tet.2024.133836.
11. Sheldon, R. A. Metrics of Green Chemistry and Sustainability: Past, Present, and Future. ACS Sustainable Chemistry & Engineering 2018, 6, 32-48. DOI: 10.1021/acssuschemeng.7b03505.
12. Hine, J.; Craig, J. C., Jr.; Underwood, J. G., II; Via, F. A. Kinetics and mechanism of the hydrolysis of N-isobutylidenemethylamine in aqueous solution. Journal of the American Chemical Society 1970, 92, 5194-5199. DOI: 10.1021/ja00720a032.
13. Hine, J.; Cholod, M. S.; Chess, W. K. Kinetics of the formation of imines from acetone and primary amines. Evidence for internal acid-catalyzed dehydration of certain intermediate carbinolamines. Journal of the American Chemical Society 1973, 95, 4270-4276. DOI: 10.1021/ja00794a025.
14. Hupe, D. J. Enzyme reactions involving imine formation. New Comprehensive Biochemistry: The Chemistry of Enzyme Action 1984, 6, 271-301. DOI: 10.1016/S0167-7306(08)60379-8.
15. Baymak, M. S.; Zuman, P. Equilibria of formation and dehydration of the carbinolamine intermediate in the reaction of benzaldehyde with hydrazine. Tetrahedron 2007, 63, 5450-5454. DOI: 10.1016/j.tet.2007.04.034.
16. Abrams, W. R.; Kallen, R. G. Equilibria and kinetics of N-hydroxymethylamine formation from aromatic exocyclic amines and formaldehyde. Effects of nucleophilicity and catalyst strength upon mechanisms of catalysis of carbinolamine formation. Journal of the American Chemical Society 1976, 98, 7777-7789. DOI: 10.1021/ja00440a052.
17. Brighente, I. M. C.; Vottero, L. R.; Terezani, A. J.; Yunes, R. A. Addition of hydroxylamine to cyclohexanone and bicyclic ketones. Steric, electronic and hydrogen bonding effects on the general mechanism of addition of amines to carbonyl compounds. Journal of Physical Organic Chemistry 1991, 4, 107-112. DOI: 10.1002/poc.610040207.
18. Rufino, V. C.; Pliego, J. R. Mechanisms of the formation of imines in aqueous solution and the effect of the pH: a theoretical analysis. Arkivoc 2020, 2020, 34-52. DOI: 10.24820/ark.5550190.p011.105.
19. Ren, X.; Liu, K.; Lu, Y.; Ding, W.; Lei, S.; Yang, L. Stepwise versus Concerted: Theoretical Insights into the Stereoselectivity in Aryl Imine Formation Assisted by Acid and Water. The Journal of Physical Chemistry A 2023, 127, 9748-9759. DOI: 10.1021/acs.jpca.3c05530.
20. Gokhale, M. Y.; Kirsch, L. E. Glycosylation of aromatic amines II: Kinetics and mechanisms of the hydrolytic reaction between kynurenine and glucose. Journal of Pharmaceutical Sciences 2009, 98, 4616-4628. DOI: 10.1002/jps.21754.
21. Kulchat, S.; Chaur, M. N.; Lehn, J.-M. Kinetic Selectivity and Thermodynamic Features of Competitive Imine Formation in Dynamic Covalent Chemistry. Chemistry - A European Journal 2017, 23, 11108-11118. DOI: 10.1002/chem.201702088.
22. Kundu, A.; Shakil, N. A.; Saxena, D. B.; Pankaj; Kumar, J.; Walia, S. Microwave assisted solvent-free synthesis and biological activities of novel imines (Schiff bases). Journal of Environmental Science and Health, Part B 2009, 44, 428-434. DOI: 10.1080/03601230902934645.
23. Eftekhari, S.; Foroughifar, N.; Hallajian, S.; Khajeh-Amiri, A. Green Synthesis of Some Novel Imidazole Schiff Base Derivatives Under Microwave Irradiation / Reflux Conditions and Evaluations of the Antibacterial Activity. Current Microwave Chemistry 2020, 7, 207-215. DOI: 10.2174/2213335607999200520124245.
24. Omar Ali, N. H. S.; Abdul Hamid, M. H. S.; Mohamad Ali Putra, N. A.; Adol, H. A.; Mirza, A. H.; Usman, A.; Siddiquee, T. A.; Hoq, M. R.; Karim, M. R. Efficient eco-friendly syntheses of dithiocarbazates and thiosemicarbazones. Green Chemistry Letters and Reviews 2020, 13, 129-140. DOI: 10.1080/17518253.2020.1737252.
25. Bhagat, S.; Sharma, N.; Chundawat, T. S. Synthesis of Some Salicylaldehyde-Based Schiff Bases in Aqueous Media. Journal of Chemistry 2013, 2013, 909217. DOI: 10.1155/2013/909217.

26. [26] Sharma, D.; Bhardwaj, A. A comparative view over the synthesis of Schiff base ligands and metal complexes by conventional and solvent free routes. International Journal of Engineering Technologies and Management Research 2020, 4, 107-117. DOI: 10.29121/ijetmr.v4.i12.2017.603.
27. Kanaiya, P. H.; Jain, V. K.; Sharma, M. G. Microwave-assisted Synthesis and Characterization of Calix[4]resorcinarene Schiff Base [C4RSB]. Current Green Chemistry 2026, 13, 302-309. DOI: 10.2174/0122133461391375250728203247.
28. Kumar, D.; Karati, D.; Mahadik, K. R. Microwave-assisted Synthetic Scheme of Novel Schiff Base Congeners of Pyrimidine Nuclei by Using Water as Solvent: A Green Approach to Synthesis. Current Microwave Chemistry 2022, 9, 3-9. DOI: 10.2174/2213335609666220414141731.
29. [Zhimomi, B. K.; Yanthan, K.; Phom, M.; Achumi, P.; Imchen, P.; Tumtin, S.; Phucho, T. Catalyst Free, Microwave-assisted Synthesis of Benzylidene Schiff Bases and Evaluation of Their Biological Properties. ChemistrySelect 2024, 9, e202404710. DOI: 10.1002/slct.202404710.
30. Manjare, S. B.; Mahadik, R. K.; Manval, K. S.; More, P. P.; Dalvi, S. S. Microwave-Assisted Rapid and Green Synthesis of Schiff Bases Using Cashew Shell Extract as a Natural Acid Catalyst. ACS Omega 2023, 8, 473-479. DOI: 10.1021/acsomega.2c05187.
31. Chakraborty, M.; Baweja, S.; Bhagat, S.; Chundawat, T. S. Microwave Assisted Synthesis of Schiff Bases: A Green Approach. International Journal of Chemical Reactor Engineering 2012, 10, A93. DOI: 10.1515/1542-6580.2973.
32. Hassan, A. M.; Said, A. O.; Heakal, B. H.; Younis, A.; Aboulthana, W. M.; Mady, M. F. Green Synthesis, Characterization, Antimicrobial and Anticancer Screening of New Metal Complexes Incorporating Schiff Base. ACS Omega 2022, 7, 32418-32431. DOI: 10.1021/acsomega.2c03911.
33. Sharma, S.; Yadav, P.; Seema; Kumari, S.; Ranka, M. Microwave-assisted synthesis of Schiff base and mixed ligand complexes of Cr(III): comparison with conventional method and antimicrobial studies. RASAYAN Journal of Chemistry 2023, 16, 884-891. DOI: 10.31788/rjc.2023.1628228.
34. Sangare, M. M.; Sankpal, S. T.; Surve, R. R. Advancements in Microwave-Assisted Synthesis of Schiff Bases and Their Metal Complexes: A Green and Efficient Approach: A Review. ChemSci Advances 2025, 2, 200-208. DOI: 10.69626/csa.2025.0200.
35. Jain, A.; De, S.; Barman, P. Microwave-assisted synthesis and notable applications of Schiff-base and metal complexes: a comparative study. Research on Chemical Intermediates 2022, 48, 2199-2251. DOI: 10.1007/s11164-022-04708-7.
36. Arafa, W. A. A.; Shaker, R. M. A facile green chemistry approaches towards the synthesis of bis-Schiff bases using ultrasound versus microwave and conventional method without catalyst. Arkivoc 2016, 2016, 187-201. DOI: 10.3998/ark.5550190.p009.464.
37. Katariya, D.; Vyas, A.; Khunt, R. Ultrasound-assisted Synthesis of Schiff Bases from 4-Aminoantipyrine: A Green and Efficient Approach. Current Science, Engineering and Technology 2026, 5, 223-228. DOI: 10.2174/0130506115401716251125061528.
38. Kargar, H.; Behjatmanesh-Ardakani, R.; Fallah-Mehrjardi, M.; Torabi, V.; Munawar, K. S.; Ashfaq, M.; Tahir, M. N. Ultrasound-based synthesis, SC-XRD, NMR, DFT, HSA of new Schiff bases derived from 2-aminopyridine: Experimental and theoretical studies. Journal of Molecular Structure 2021, 1233, 130105. DOI: 10.1016/j.molstruc.2021.130105.
39. Mousavi, S. A.; Montazerozohori, M.; Masoudiasl, A.; Mahmoudi, G.; White, J. M. Sonication-assisted synthesis of a new cationic zinc nitrate complex with a tetradentate Schiff base ligand: Crystal structure, Hirshfeld surface analysis and investigation of different parameters influence on morphological properties. Ultrasonics Sonochemistry 2018, 46, 26-35. DOI: 10.1016/j.ultsonch.2018.02.050.
40. Zare, N.; Zabardasti, A. A new nano-sized mononuclear Cu(II) complex with N,N-donor Schiff base ligands: sonochemical synthesis, characterization, molecular modeling and biological activity. Applied Organometallic Chemistry 2019, 33, e4687. DOI: 10.1002/aoc.4687.
41. Abdel-Rahman, L. H.; Abu-Dief, A. M.; El-Khatib, R. M.; Abdel-Fatah, S. M. Some new nano-sized Fe(II), Cd(II) and Zn(II) Schiff base complexes as precursor for metal oxides: Sonochemical synthesis, characterization, DNA interaction, in vitro antimicrobial and anticancer activities. Bioorganic Chemistry 2016, 69, 140-152. DOI: 10.1016/j.bioorg.2016.10.009.
42. Saleh, A. M.; Mahdy, A. G.; Hamed, A. A. Sonochemically synthesized Ag(I) and Ni(II) Schiff base complexes as efficient visible-light photocatalysts for dye degradation with DFT insights. Scientific Reports 2026, 16, 7181. DOI: 10.1038/s41598-026-37498-8.
43. Arami, J. A.; Yang, X.; Brodie, A. W.; McElwee-White, L.; Yap, G. P. A.; Li, X.; Jurca, T. One-Pot, One-Step Mn-bis(imino)pyridine Complexes through Sonochemistry. Inorganic Chemistry 2026, 65, 16639-16644. DOI: 10.1021/acs.inorgchem.6c01196.
44. Sanjurani, T.; Paul, S.; Barman, P. Indole-based NNN donor Schiff base ligand and its complexes: Sonication-assisted synthesis, characterization, DNA binding, anti-cancer evaluation and in-vitro biological assay. Bioorganic Chemistry 2024, 146, 107281. DOI: 10.1016/j.bioorg.2024.107281.

45. Elseman, A. M.; Shalan, A. E.; Rashad, M. M.; Hassan, A. M.; Ibrahim, N. M.; Nassar, A. M. Easily attainable new approach to mass yield ferrocenyl Schiff base and different metal complexes of ferrocenyl Schiff base through convenient ultrasonication-solvothermal method. Journal of Physical Organic Chemistry 2017, 30, e3639. DOI: 10.1002/poc.3639.
46. Friscic, T.; Mottillo, C.; Titi, H. M. Mechanochemistry for Synthesis. AngewandteChemie International Edition 2020, 59, 1018-1029. DOI: 10.1002/anie.201906755.
47. Wang, G.-W. Mechanochemical organic synthesis. Chemical Society Reviews 2013, 42, 7668-7700. DOI: 10.1039/C3CS35526H.
48. Ciesielska, K.; Hoffmann, M.; Kubicki, M.; Pluskota-Karwatka, D. Mechanochemical Synthesis of Fluorinated Imines. Molecules 2022, 27, 4557. DOI: 10.3390/molecules27144557.
49. Rashmi, P. S.; Ramakrishna, D.; Supriya, S.; Ranganatha, S. Mechanochemistry in Action: Efficient, Solvent-free Synthesis of Chromone-Derived Imine Compounds and Their Biological Screening. ChemistrySelect 2026, 11, e202504966. DOI: 10.1002/slct.202504966.
50. Ohn, N.; Kim, J. G. Mechanochemical Post-Polymerization Modification: Solvent-Free Solid-State Synthesis of Functional Polymers. ACS Macro Letters 2018, 7, 561-565. DOI: 10.1021/acsmacrolett.8b00171.
51. Fantozzi, N.; Volle, J.-N.; Porcheddu, A.; Virieux, D.; Garcia, F.; Colacino, E. Green metrics in mechanochemistry. Chemical Society Reviews 2023, 52, 6680-6714. DOI: 10.1039/D2CS00997H.
52. Pattanayak, P.; Saha, S.; Chatterjee, T.; Ranu, B. C. Sustainable and solvent-free synthesis of molecules of pharmaceutical importance by ball milling. Chemical Communications 2025, 61, 247-265. DOI: 10.1039/D4CC05127K.
53. Komar, M.; Prasnikar, F.; Gazivoda Kraljevic, T.; Aladic, K.; Molnar, M. 3-Amino-2-methylquinazolin-4-(3H)-one Schiff bases synthesis - a green chemistry approach - a comparison of microwave and ultrasound promoted synthesis with mechanosynthesis. Current Green Chemistry 2021, 8, 62-69. DOI: 10.2174/2213346107999201231125434.
54. Patil, S.; Jadhav, S. D.; Deshmukh, M. B.; Patil, U. P. Natural acid catalyzed synthesis of Schiff base under solvent-free condition: as a green approach. International Journal of Organic Chemistry 2012, 2, 166-171. DOI: 10.4236/ijoc.2012.22025.
55. Ma'rufah, L.; Hanapi, A.; Ningsih, R.; Fasya, A. G. Synthesis of Schiff Base Compounds from Vanillin and p-Aminoacetophenone Using Lime Juice as a Natural Acid Catalyst and Their Utilization as Corrosion Inhibitors. Advances in Social Science, Education and Humanities Research 2021, 529, 253-258. DOI: 10.2991/assehr.k.210421.043.
56. Sachdeva, H.; Saroj, R.; Khaturia, S.; Dwivedi, D.; Chauhan, O. P. Green Route for Efficient Synthesis of Novel Amino Acid Schiff Bases as Potent Antibacterial and Antifungal Agents and Evaluation of Cytotoxic Effects. Journal of Chemistry 2014, 2014, 848543. DOI: 10.1155/2014/848543.
57. Bentoumi, H.; Tliba, S.; K'tir, H.; Chohra, D.; Aouf, Z.; Adjeroud, Y.; Amira, A.; Zerrouki, R.; Ibrahim-Ouali, M.; Aouf, N.-E.; Liacha, M. Experimental synthesis, biological evaluation, theoretical investigations of some novel benzoxazolinone based Schiff bases under eco-environmental conditions as potential antioxidant agents. Journal of Molecular Structure 2022, 1270, 133986. DOI: 10.1016/j.molstruc.2022.133986.
58. Shaik, B. B.; Seboletswe, P.; Mohite, S. B.; Katari, N. K.; Bala, M. D.; Karpoormath, R.; Singh, P. Lemon Juice: A Versatile Biocatalyst and Green Solvent in Organic Transformations. ChemistrySelect 2022, 7, e202103701. DOI: 10.1002/slct.202103701.
59. [Nigam, M.; Tuttle, D.; Morra, B.; Dicks, A. P.; Rodriguez, J. Putting the squeeze on imine synthesis: citrus juice as a reaction medium in the introductory organic laboratory. Green Chemistry Letters and Reviews 2023, 16, 2185107. DOI: 10.1080/17518253.2023.2185107.
60. Bhusari, N.; Bagul, A.; Mishra, V. K.; Tufail, A.; Gaikwad, D.; Dubey, A. Sustainable synthesis of Schiff base derivatives via an ionic liquid and a microwave-assisted approach: structural, biological, and computational evaluation. RSC Advances 2025, 15, 22764-22788. DOI: 10.1039/D5RA02622A.
61. Zhang, Q.; De Oliveira Vigier, K.; Royer, S.; Jerome, F. Deep eutectic solvents: syntheses, properties and applications. Chemical Society Reviews 2012, 41, 7108-7146. DOI: 10.1039/C2CS35178A.
62. Alonso, D. A.; Baeza, A.; Chinchilla, R.; Guillena, G.; Pastor, I. M.; Ramon, D. J. Deep Eutectic Solvents: The Organic Reaction Medium of the Century. European Journal of Organic Chemistry 2016, 2016, 612-632. DOI: 10.1002/ejoc.201501197.
63. Perna, F. M.; Vitale, P.; Capriati, V. Deep eutectic solvents and their applications as green solvents. Current Opinion in Green and Sustainable Chemistry 2020, 21, 27-33. DOI: 10.1016/j.cogsc.2019.09.004.
64. Matsumoto, M.; Dasari, R. R.; Ji, W.; Feriante, C. H.; Parker, T. C.; Marder, S. R.; Dichtel, W. R. Rapid, Low Temperature Formation of Imine-Linked Covalent Organic Frameworks Catalyzed by Metal Triflates. Journal of the American Chemical Society 2017, 139, 4999-5002. DOI: 10.1021/jacs.7b01240.
65. Tamura, M.; Tomishige, K. Scope and reaction mechanism of CeO2-catalyzed one-pot imine synthesis from alcohols and amines. Journal of Catalysis 2020, 389, 285-296. DOI: 10.1016/j.jcat.2020.05.031.

66. Lu, X.; Zhang, F.; Yan, F.; Liu, Q.; Zhao, Y.; Du, M.; Zhang, L.; Li, X.; Zhao, Z.; Liu, H. Pd-catalyzed imine synthesis from nitroarenes via Fe-H2O under electromagnetic milling. Chemical Communications 2025, 61, 6316-6319. DOI: 10.1039/D5CC00498E.
67. Banerjee, S.; Periyasamy, S.; Muthukumaradoss, K.; Deivasigamani, P.; Saravanan, V. Revolutionizing organic synthesis through green chemistry: metal-free, bio-based, and microwave-assisted methods. Frontiers in Chemistry 2025, 13, 1656935. DOI: 10.3389/fchem.2025.1656935.
68. Martin-Illan, J. A.; Rodriguez-San-Miguel, D.; Franco, C.; Imaz, I.; Maspoch, D.; Puigmarti-Luis, J.; Zamora, F. Green synthesis of imine-based covalent organic frameworks in water. Chemical Communications 2020, 56, 6704-6707. DOI: 10.1039/D0CC02033H.
69. Wang, X.-L.; Zhang, L.-T.; He, S.; Chen, X.-X.; Huang, X.-C.; Zhou, H.-L. Dynamic Imine Exchange Reactions for Facile Synthesis of Imine-Linked Covalent Organic Frameworks. Chemistry of Materials 2023, 35, 10070-10077. DOI: 10.1021/acs.chemmater.3c02092.
70. Maleki, A.; Javanshir, S.; Naimabadi, M. Facile synthesis of imidazo[1,2-a]pyridines via a one-pot three-component reaction under solvent-free mechanochemical ball-milling conditions. RSC Advances 2014, 4, 30229-30232. DOI: 10.1039/C3RA43221A.
71. Naeimi, H.; Nazifi, Z. S. Efficient and convenient template synthesis and characterization of copper(II) and cobalt(II) complexes of 1,1'-[arenediylbis(nitrilomethylidine)]bis-2-naphtholes. Chemija 2013, 24, 191-196. DOI: 10.6001/chemija.2013.24.3.4.
72. Nelson, S. M. Developments in the synthesis and coordination chemistry of macrocyclic Schiff base ligands. Pure and Applied Chemistry 1980, 52, 2461-2476. DOI: 10.1351/pac198052112461.
73. Radecka-Paryzek, W.; Patroniak, V.; Lisowski, J. Metal complexes of polyaza and polyoxaaza Schiff base macrocycles. Coordination Chemistry Reviews 2005, 249, 2156-2175. DOI: 10.1016/j.ccr.2005.02.021.
74. Zhao, P.-C.; Chang, F.-F.; Feng, F.-D.; Huang, W. Transmetalation and Demetallization for Open-Oyster-like Non-Ionic Cd(II) Macrocycles. Inorganic Chemistry 2020, 59, 7504-7511. DOI: 10.1021/acs.inorgchem.0c00304.
75. Yan, S.; Feng, G.; Geng, J.; Feng, F.; Ma, H.; Huang, W. Tunable Construction of Sandwich-Type Double-[1+1] and Half-Folded [2+2] Schiff-Base Complexes Controlled by the Combination of Primary and Secondary Template Effects. Inorganic Chemistry 2022, 61, 20994-21003. DOI: 10.1021/acs.inorgchem.2c03473.
76. Pandey, P.; Verma, A.; Bretosh, K.; Sutter, J.-P.; Sunkari, S. S. Template directed synthesis of half condensed Schiff base complexes of Cu(II) and Co(III): Structural and magnetic studies. Polyhedron 2019, 164, 80-89. DOI: 10.1016/j.poly.2019.02.037.
77. Mazzoni, R.; Roncaglia, F.; Rigamonti, L. When the Metal Makes the Difference: Template Syntheses of Tridentate and Tetradentate Salen-Type Schiff Base Ligands and Related Complexes. Crystals 2021, 11, 483. DOI: 10.3390/cryst11050483.
78. Rezaeivala, M.; Keypour, H. Schiff base and non-Schiff base macrocyclic ligands and complexes incorporating the pyridine moiety - The first 50 years. Coordination Chemistry Reviews 2014, 280, 203-253. DOI: 10.1016/j.ccr.2014.06.007.
79. Remiya, J. P.; Sikha, T. S.; Shyni, B. One-pot synthesis and characterization of Schiff base macrocyclic complexes as a potential bioactive core - a review. Journal of Coordination Chemistry 2021, 74, 3081-3108. DOI: 10.1080/00958972.2021.2025223.
80. Lisowski, J. Imine- and Amine-Type Macrocycles Derived from Chiral Diamines and Aromatic Dialdehydes. Molecules 2022, 27, 4097. DOI: 10.3390/molecules27134097.
81. Rajakkani, P.; Alagarraj, A.; GurusamyThangavelu, S. A. Tetraaza macrocyclic Schiff base metal complexes bearing pendant groups: Synthesis, characterization and bioactivity studies. Inorganic Chemistry Communications 2021, 134, 108989. DOI: 10.1016/j.inoche.2021.108989.
82. Shailakshi; Tees, R.; D'souza, G.; Sapna Kumari, M.; Pinto, G. M.; Priya A Comparative Study of Green and Conventional Synthesis of Schiff Base and its Metal Complexes from 3,5-Dichloroaniline and 3-Chloro-4-fluoroaniline. International Research Journal on Advanced Engineering and Management 2024, 2, 2323-2327. DOI: 10.47392/irjaem.2024.0336.
83. Sharma, A.; Arora, S. K. A review study on green synthesis of Schiff bases. Indian Journal of Applied Research 2023, 13, 69-72. DOI: 10.36106/ijar/9529205.
84. Roman, G. Mannich bases in medicinal chemistry and drug design. European Journal of Medicinal Chemistry 2015, 89, 743-816. DOI: 10.1016/j.ejmech.2014.10.076.
85. Karthikeyan, M. S.; Prasad, D. J.; Poojary, B.; Bhat, K. S.; Holla, B. S.; Kumari, N. S. Synthesis and biological activity of Schiff and Mannich bases bearing 2,4-dichloro-5-fluorophenyl moiety. Bioorganic & Medicinal Chemistry 2006, 14, 7482-7489. DOI: 10.1016/j.bmc.2006.07.015.
86. Csuvik, O.; Szatmari, I. Synthesis of Bioactive Aminomethylated 8-Hydroxyquinolines via the Modified Mannich Reaction. International Journal of Molecular Sciences 2023, 24, 7915. DOI: 10.3390/ijms24097915.
87. Raoof, S. S.; Sadiq, A. S. Mannich bases: synthesis, pharmacological activity, and applications: a review. Iraqi Journal of Science 2022, 63, 5086-5105. DOI: 10.24996/ijs.2022.63.12.1.

88. Benkhaya, S.; M'rabet, S.; El Harfi, A. Classifications, properties, recent synthesis and applications of azo dyes. Heliyon 2020, 6, e03271. DOI: 10.1016/j.heliyon.2020.e03271.
89. Khanum, R.; Shoukat Ali, R. A.; Santosh Kumar, S. R.; Husnasarvari, M.; Rangaswamy, H. R. Synthesis, characterization and versatile applications of 1,4-benzodioxan-6-amine based novel azo dyes. Inorganic Chemistry Communications 2023, 157, 111297. DOI: 10.1016/j.inoche.2023.111297.
90. Al-Haj Hussien, F. An eco-friendly methodology for the synthesis of azocoumarin dye using cation exchange resins. Heliyon 2021, 7, e08439. DOI: 10.1016/j.heliyon.2021.e08439.
91. Khushi Siri, C. N.; Shoukat Ali, R. A.; Latha, M. S.; Betageri, V. S.; Byadagi, K. S. A reflection of literature reports on synthesis, spectral studies, multilateral applications of azo dyes. Results in Chemistry 2025, 18, 102878. DOI: 10.1016/j.rechem.2025.102878.
92. Gomez, S.; Peters, J. A.; Maschmeyer, T. The Reductive Amination of Aldehydes and Ketones and the Hydrogenation of Nitriles: Mechanistic Aspects and Selectivity Control. Advanced Synthesis & Catalysis 2002, 344, 1037-1057. DOI: 10.1002/1615-4169(200212)344:10<1037::AID-ADSC1037>3.0.CO;2-3.
93. Irrgang, T.; Kempe, R. Transition-Metal-Catalyzed Reductive Amination Employing Hydrogen. Chemical Reviews 2020, 120, 9583-9674. DOI: 10.1021/acs.chemrev.0c00248.
94. Murugesan, K.; Senthamarai, T.; Chandrashekhar, V. G.; Natte, K.; Kamer, P. C. J.; Beller, M.; Jagadeesh, R. V. Catalytic reductive aminations using molecular hydrogen for synthesis of different kinds of amines. Chemical Society Reviews 2020, 49, 6273-6328. DOI: 10.1039/C9CS00286C.
95. Aleku, G. A.; France, S. P.; Man, H.; Mangas-Sanchez, J.; Montgomery, S. L.; Sharma, M.; Leipold, F.; Hussain, S.; Grogan, G.; Turner, N. J. A reductive aminase from Aspergillus oryzae. Nature Chemistry 2017, 9, 961-969. DOI: 10.1038/nchem.2782.
96. Meldal, M.; Tornoe, C. W. Cu-Catalyzed Azide-Alkyne Cycloaddition. Chemical Reviews 2008, 108, 2952-3015. DOI: 10.1021/cr0783479.
97. Bock, V. D.; Hiemstra, H.; van Maarseveen, J. H. CuI-Catalyzed Alkyne-Azide "Click" Cycloadditions from a Mechanistic and Synthetic Perspective. European Journal of Organic Chemistry 2006, 2006, 51-68. DOI: 10.1002/ejoc.200500483.
98. Aflak, N.; Ben El Ayouchia, H.; Bahsis, L.; Anane, H.; Julve, M.; Stiriba, S.-E. Recent Advances in Copper-Based Solid Heterogeneous Catalysts for Azide-Alkyne Cycloaddition Reactions. International Journal of Molecular Sciences 2022, 23, 2383. DOI: 10.3390/ijms23042383.
99. Meng, G.; Guo, T.; Ma, T.; Zhang, J.; Shen, Y.; Sharpless, K. B.; Dong, J. Modular click chemistry libraries for functional screens using a diazotizing reagent. Nature 2019, 574, 86-89. DOI: 10.1038/s41586-019-1589-1.
100. Kaur, J.; Saxena, M.; Rishi, N. An Overview of Recent Advances in Biomedical Applications of Click Chemistry. Bioconjugate Chemistry 2021, 32, 1455-1471. DOI: 10.1021/acs.bioconjchem.1c00247.
101. Sheldon, R. A. The E factor 25 years on: the rise of green chemistry and sustainability. Green Chemistry 2017, 19, 18-43. DOI: 10.1039/C6GC02157C.
102. Ribeiro, M. G. T. C.; Machado, A. A. S. C. Greenness of chemical reactions - limitations of mass metrics. Green Chemistry Letters and Reviews 2013, 6, 1-18. DOI: 10.1080/17518253.2012.669798.
103. Kerek, A. L.; Sens, L.; Sandri, M. C. M.; Fiorin, B. C. Green synthesis of azachalcones: an evaluation based on chemical greenness metrics. Green Chemistry Letters and Reviews 2025, 18, 2581930. DOI: 10.1080/17518253.2025.2581930.
104. Fotopoulou, E.; Anastasiou, P. K.; Tomza, C.; Neochoritis, C. G. The Ugi reaction as the green alternative towards active pharmaceutical ingredients. Tetrahedron Green Chemistry 2024, 3, 100044. DOI: 10.1016/j.tgchem.2024.100044.
105. Pushpanathan, V.; Dhas, S. S. J.; Kumar, D. S. Synthesis, spectroscopic and surface morphological investigations of Co(II), Ni(II), Cu(II), and Zn(II) Schiff base macrocyclic complexes. Chemical Physics Impact 2023, 7, 100264. DOI: 10.1016/j.chphi.2023.100264.
106. Qian, C.; Feng, L.; Teo, W. L.; Liu, J.; Zhou, W.; Wang, D.; Zhao, Y. Imine and imine-derived linkages in two-dimensional covalent organic frameworks. Nature Reviews Chemistry 2022, 6, 881-898. DOI: 10.1038/s41570-022-00437-y.
107. Alharthi, N. H.; El-Hashemy, M. A.; Derafa, W. M.; Althobaiti, I. O.; Altaleb, H. A. Corrosion inhibition of mild steel by highly stable polydentate Schiff base derived from 1,3-propanediamine in aqueous acidic solution. Journal of Saudi Chemical Society 2022, 26, 101501. DOI: 10.1016/j.jscs.2022.101501.
108. Rep Kaulic, V.; PantalonJuraj, N.; Peric, B.; Stefanic, Z.; Kirin, S. I.; Raic-Malic, S. Transition Metal-Mediated Hydrolysis of the Imine Bond in 2-Azomethine Benzothiazoles. CroaticaChemica Acta 2023, 95, 123-130. DOI: 10.5562/cca3907.
109. Halder, A.; Karak, S.; Addicoat, M.; Bera, S.; Chakraborty, A.; Kunjattu, S. H.; Pachfule, P.; Heine, T.; Banerjee, R. Ultrastable Imine-Based Covalent Organic Frameworks for Sulfuric Acid Recovery: An Effect of Interlayer Hydrogen Bonding. AngewandteChemie International Edition 2018, 57, 5797-5802. DOI: 10.1002/anie.201802220.

110. Cusin, L.; Peng, H.; Ciesielski, A.; Samori, P. Chemical Conversion and Locking of the Imine Linkage: Enhancing the Functionality of Covalent Organic Frameworks. AngewandteChemie International Edition 2021, 60, 14236-14250. DOI: 10.1002/anie.202016667.
111. Li, X.; Zhang, C.; Cai, S.; Lei, X.; Altoe, V.; Hong, F.; Urban, J. J.; Ciston, J.; Chan, E. M.; Liu, Y. Facile transformation of imine covalent organic frameworks into ultrastable crystalline porous aromatic frameworks. Nature Communications 2018, 9, 2998. DOI: 10.1038/s41467-018-05462-4.
112. Anjana, P. N.; Pulikkal, A. K. Synthesis, derivation, and applications of imine-linked covalent organic frameworks: A comprehensive review. Microporous and Mesoporous Materials 2025, 387, 113516. DOI: 10.1016/j.micromeso.2025.113516.
113. Gupta, S. K.; Mitra, R. K.; Yadav, M.; Dagdag, O.; Berisha, A.; Mamba, B. B.; Nkambule, T. T. I.; Ebenso, E. E.; Singh, S. K. Electrochemical, surface morphological and computational evaluation on carbohydrazide Schiff bases as corrosion inhibitor for mild steel in acidic medium. Scientific Reports 2023, 13, 15108. DOI: 10.1038/s41598-023-41975-9.
114. Okey, N. C.; Obasi, N. L.; Ejikeme, P. M.; Ndinteh, D. T.; Ramasami, P.; Sherif, E.-S. M.; Akpan, E. D.; Ebenso, E. E. Evaluation of some amino benzoic acid and 4-aminoantipyrine derived Schiff bases as corrosion inhibitors for mild steel in acidic medium: Synthesis, experimental and computational studies. Journal of Molecular Liquids 2020, 315, 113773. DOI: 10.1016/j.molliq.2020.113773.
115. Chugh, B.; Singh, A. K.; Thakur, S.; Pani, B.; Lgaz, H.; Chung, I.-M.; Jha, R.; Ebenso, E. E. Comparative Investigation of Corrosion-Mitigating Behavior of Thiadiazole-Derived Bis-Schiff Bases for Mild Steel in Acid Medium: Experimental, Theoretical, and Surface Study. ACS Omega 2020, 5, 13503-13520. DOI: 10.1021/acsomega.9b04274.
116. El Guesmi, N.; Asghar, B. H.; Awad, M. I.; Al Harbi, A. N.; Kassem, M. A.; Shaaban, M. R. Novel thiazole-derived Schiff bases as efficient corrosion inhibitors for mild steel in acidic media: Synthesis, electrochemical and computational insights. Arabian Journal of Chemistry 2024, 17, 105867. DOI: 10.1016/j.arabjc.2024.105867.
117. Menaka, R.; Subhashini, S. Chitosan Schiff base as effective corrosion inhibitor for mild steel in acid medium. Polymer International 2017, 66, 349-358. DOI: 10.1002/pi.5245.
118. Afrin, A.; Chinna Ayya Swamy, P. Novel Schiff base derivatives for the detection of one-to-multi metal ions and tracking the live cell imaging. Coordination Chemistry Reviews 2023, 494, 215327. DOI: 10.1016/j.ccr.2023.215327.
119. Musikavanhu, B.; Liang, Y.; Xue, Z.; Feng, L.; Zhao, L. Strategies for Improving Selectivity and Sensitivity of Schiff Base Fluorescent Chemosensors for Toxic and Heavy Metals. Molecules 2023, 28, 6960. DOI: 10.3390/molecules28196960.
120. Jacobsen, E. N. Asymmetric Catalysis of Epoxide Ring-Opening Reactions. Accounts of Chemical Research 2000, 33, 421-431. DOI: 10.1021/ar960061v.
121. Canali, L.; Sherrington, D. C. Utilisation of homogeneous and supported chiral metal(salen) complexes in asymmetric catalysis. Chemical Society Reviews 1999, 28, 85-93. DOI: 10.1039/A806483K.